

Critical Issues and Trends: Underserved Populations; Fitness

The Conspicuous Absence of People With Disabilities in Public Fitness and Recreation Facilities: Lack of Interest or Lack of Access?

James H. Rimmer, PhD



INTRODUCTION

The lack of participation in beneficial exercise is a serious public health concern for all Americans, but it is even more acute for the estimated 52 million Americans with disabilities who are demonstrably at much greater risk for developing the types of serious health problems associated with a sedentary lifestyle.¹⁻⁶ Despite the enormous health benefits that can be derived from regular physical activity,⁷ people with disabilities remain one of the most physically inactive groups in society.⁸⁻¹⁰ *Healthy People 2010* uses cross-sectional surveys to outline current levels of physical activity and exercise for various subpopulations in the United States and lists goals for the year 2010.⁶ As shown in Table 1, people with disabilities are currently much less active than their nondisabled counterparts and participate in less regular and vigorous physical activity. They also report a substantially high number of secondary conditions, which are directly or indirectly associated with their disability but in most cases are considered preventable (e.g., fatigue, weight gain, pain).¹¹

Efforts to eliminate health disparities must address issues, needs, and barriers of people with disabilities for positive lifestyle change. The level of physical inactivity observed among people with disabilities has been linked to an increase in the severity of disability and erosion of in-

volvement in community activities.¹² These patterns of low physical activity reported among people with disabilities raise serious concern regarding their health and well being, particularly as they enter their later years when the effects of the natural aging process are compounded by years of sedentary living and severe deconditioning.¹³⁻¹⁵

DISABILITY-UNFRIENDLY PHYSICAL ACTIVITY ENVIRONMENTS

The *Healthy People 2010* chapter entitled “Disability and Secondary Conditions” notes that the significantly lower rate of participation among people with disabilities in physical activity may be related to environmental barriers—including architectural barriers, organizational policies and practices, discrimination, and social attitudes—and recommends that public health agencies begin to evaluate which environmental factors enhance or impede participation in physical activity.⁶ Presently, many people with disabilities consider most local and state parks, fitness centers, health clubs, spas, gymnasiums, playgrounds, pools, trails, and sports fields to be “unfriendly” environments because of a lack of accessibility.¹⁶⁻¹⁸

Physical environments, both indoor and outdoor, have an important role in influencing participation in physical activity.¹⁹⁻²² Factors that affect accessibility include the built environment, equipment, information, staff training, and policies and procedures.

Physical (Built) Environment

Indoor environments of many fitness facilities and health clubs impose substantial barriers to participation among people with disabilities.^{17,18,22} Doors are often difficult to open; bathrooms are not always accessible, particularly in smaller gymnasiums and health clubs; and stairs often lead from one part of the facility to another (e.g., cardiovascular equipment is on one floor and strength equipment is on another floor). Although the Americans with Disabilities Act requires facilities to be architecturally accessible, many facilities are not in compliance with the federal law.²³

James H. Rimmer, PhD, is a Professor in the Department of Disability and Human Development, University of Illinois at Chicago.

Send reprint requests to James H. Rimmer, PhD, Director, National Center on Physical Activity and Disability, and Director, Rehabilitation Engineering Research Center—RecTech, Department of Disability and Human Development, University of Illinois at Chicago, 1640 West Roosevelt Road, Chicago, IL 60608-6904; jrimmer@uic.edu.

This manuscript was submitted March 30, 2004; revisions were requested August 24, 2004, and November 10, 2004; the manuscript was accepted for publication November 17, 2004.

*Am J Health Promot 2005;19(5):327-329.
Copyright © 2005 by American Journal of Health Promotion, Inc.
0890-1171/05/\$5.00 + 0*

Table 1
Healthy People 2010
Goals for Increasing Physical Activity

	% People With Disabilities	% People Without Disabilities	% Healthy People 2010 Target
No leisure-time physical activity	56	36	20
30 min of moderate physical activity 5+ days per week	12	16	30
20 min of vigorous physical activity 3+ days per week	13	25	30

The outdoor environment can be even less accessible than indoor facilities. Streets often do not have curb cuts; sidewalks are broken and damaged, creating a higher risk of falling; walkways in parks and recreation areas are too narrow for a wheelchair user and partner walking side-by-side; the community does not have sidewalks; paths are poorly lit; many urban and suburban communities have a high traffic volume and excessive noise from motor vehicles; or the terrain is too steep. Other potential problems with outdoor environments include unsafe neighborhoods, poor weather that makes sidewalks slippery or impassable, too few benches along a trail for people who need frequent rest periods, poorly designated signage, no accessible bathrooms along a trail or path, and no handicapped parking spaces close to a trail.²⁴

Exercise Equipment

Most manufacturers of exercise equipment do not consider in their design specifications how to make their equipment accessible for people with physical, cognitive, and sensory disabilities. Inaccessible equipment is a major problem in terms of participation in fitness-enhancing activities for people with disabilities. Most commercial cardiovascular exercise equipment requires propulsion by using the musculature of both lower extremities (i.e., treadmills, stationary bikes, elliptical cross-trainers, and step-pers), thereby restricting use among people with lower extremity disabilities (e.g., paralysis, limb loss). Although a few fitness facilities may be able to purchase a commercial-quality arm cycle or wheelchair ergometer, the vast majority of fitness centers either cannot afford this equipment or do not find it cost effective to purchase one for a small percentage of their clientele.²⁵ Another problem with inaccessible equipment is that offering clients with disabilities the opportunity to use one piece of adaptive exercise equipment when the rest of the membership has access to all the equipment clearly limits the amount of enjoyment and benefit that can be obtained from a more diversified program.

People with visual or cognitive disabilities also have problems using various types of exercise equipment. Display panels are often difficult to read or understand, getting on and off the equipment presents some risk of fall-

ing, and machines are often hard to propel or lift (e.g., weights are too heavy) for people with low strength levels.

Increased access to standard exercise equipment by people with disabilities would be beneficial to individuals with physical, cognitive, and sensory limitations. Many find it difficult to use the variety of exercise machines currently available to people without disabilities to enhance cardiovascular and musculoskeletal health.

Walking Is Often the Recommended Modality for Increasing Physical Activity

Public health messages often include recommendations that encourage increased walking as the primary mode of physical activity, taking the stairs whenever possible, or parking farther away from work and walking the remaining distance. The current initiative, Steps to a Healthier U.S., sponsored by the Department of Health and Human Services, encourages people to use pedometers to walk 10,000 steps a day either through work-related activity (e.g., walking to and from work) or general activity. Unfortunately, this initiative is not targeted to people with disabilities and chronic health conditions who are unable to walk or have difficulty walking, and inexpensive and simple pedometers are unavailable to monitor physical activity in wheelchair users.

It is understandable why many public health organizations recommend increased walking as the primary mode for increasing physical activity. The majority of the population is able to walk, and walking does not involve any cost or skill and can be done at convenient times during the day (e.g., to and from work). However, for people with disabilities and chronic health conditions who have lower limb paralysis or weakness (e.g., spinal cord injury, multiple sclerosis, cerebral palsy, Parkinsons disease); arthritis; knee, back, or hip pain; extreme obesity; pulmonary diseases; or other types of disabling conditions, this recommendation is not a realistic option. Alternative forms of physical activity, such as performing various types of chair-exercise regimens that can be done at home or outdoors (e.g., chair-exercise videos or television shows, resistance exercise with milk containers or soup cans, wheeling in a local park), using upper body exercise equipment such as an arm ergometer, or having access to a swimming pool at a local public fitness facility should be recommended so that people with disabilities can participate in a parallel set of public health recommendations that achieve similar health benefits. Many YMCAs and park district programs offer discounted memberships for people with disabilities.

Commercial and Print Media

Much of the commercial and print media portray slender individuals participating in various types of physical activity. The concept of being physically active and disabled is not a common visualization in the mainstream media. The lack of photos of people with disabilities exercising outdoors or in health clubs and recreational facilities is a missed opportunity to encourage greater participation among a subgroup with some of the highest levels of physical inactivity. Additionally, brochures and other

print materials produced by private and public fitness facilities seldom provide information on what services and programs are available to people with disabilities.²⁶

PLANNING FOR A BETTER FUTURE

Physical activity could have substantial benefits in improving the health of people with disabilities by reducing the incidence of chronic diseases (e.g., type 2 diabetes, heart disease, obesity), improving secondary conditions associated with various disabilities (e.g., weakness, fatigue, reduced mobility, joint stiffness, social isolation, depression), and allowing individuals to maintain a higher level of independence in performing various activities of daily living and instrumental activities of daily living.²⁷⁻³⁰ Public and private health organizations, as well as industry and policymakers, must work together in addressing the large and growing disparity in physical activity participation observed between people with and without disabilities.⁶

Future promotional campaigns focusing on greater participation in physical activity among the U.S. population must present a more inclusive image that suggests to people with disabilities that they, too, should participate in regular physical activity most days of the week. Organizations working to increase physical activity among the general population should also use resources that provide services for people with disabilities. One example is the National Center on Physical Activity and Disability (www.ncpad.org, 1-800-900-8086), which is a federally funded information center containing many useful physical activities for people with varying types of disabilities, including home-exercise videos for upper body exercise and a list of organizations nationwide that provide disability-friendly physical activity programs. Increasing access to physical activity for the more than 50 million Americans with disabilities will take a cohesive and structured plan that emphasizes equal access for everybody.

Acknowledgments


This work was supported in part by the Centers for Disease Control and Prevention; National Center on Birth Defects and Developmental Disabilities, Disability and Health Branch R04CCR518810, U59CCU516732; and the National Institute on Disability and Rehabilitation Research H133E020715.

References

1. Binder EF, Burge SJ, Spina R. Peak aerobic power as an important component of physical performance in older women. *J Gerontol A Biol Sci Med Sci.* 1999;54:M353-M356.
2. Cooper RA, Quatrano LA, Axelson PW, et al. Research on physical activity and health among people with disabilities: a consensus statement. *J Rehabil Res Dev.* 1999;36:142-154.
3. Durstine JL, Painter PP, Franklin BA, et al. Physical activity for the chronically ill and disabled. *Sports Med.* 2000;30:207-219.
4. Lawrence RH, Jette AM. Disentangling the disablement process. *J Gerontol B Psychol Sci Soc Sci.* 1996;51:5173-5182.
5. Rimmer JH, Braddock D. Physical activity, disability, and cardiovascular health. In: Leon AS, ed. *Physical Activity and Cardiovascular Health. A National Consensus.* Champaign, Ill: Human Kinetics; 1997:236-244.
6. US Department of Health and Human Services. *Healthy People 2010.* (Conference edition in two volumes). Washington, DC: US Government Printing Office; 2000.
7. US Department of Health and Human Services. *Physical Activity and Health: A Report of the Surgeon General.* Atlanta, Ga: US Dept of Health and Human Services, Centers for Disease Control and Prevention, National Center for Chronic Disease and Prevention; 1996.
8. Finch C, Owen N, Price R. Current injury or disability as a barrier to being more physically active. *Med Sci Sports Exerc.* 2001;33:778-782.
9. Heath GW, Fentem PH. Physical activity among persons with disabilities: a public health perspective. *Exerc Sport Sci Rev.* 1997;25:195-234.
10. Nosek MA, Howland CA, Rintala DH, et al. *National Study of Women with Physical Disabilities: Final Report.* Houston, Tex: Center for Research on Women with Disabilities; 1997.
11. Kinne S, Patrick DL, Doyle DL. Prevalence of secondary conditions among people with disabilities. *Am J Public Health.* 2004;94:443-445.
12. Rimmer JH, Rubin SS, Braddock D, Hedman GH. Physical activity patterns in African-American women with physical disabilities. *Med Sci Sports Exerc.* 1999;31:613-618.
13. Ravesloot C, Seekins T, Young Q. Health promotion for people with chronic illness and physical disabilities: the connection between health psychology and disability prevention. *Clin Psychol Psychother.* 1998;5:76-85.
14. Rejeski JW, Focht B. Aging and physical disability: on integrating group and individual counseling with the promotion of physical activity. *Exerc Sport Sci Rev.* 2002;30:166-170.
15. Lexell J. Muscle structure and function in chronic neurological disorders: the potential of exercise to improve activities of daily living. *Exerc Sport Sci Rev.* 2000;28:80-84.
16. Bullock CC, Mahon MJ. Disability awareness: considerations for the exercise leader. In: Miller PD, ed. *Fitness Programming and Physical Disability.* Champaign, Ill: Human Kinetics; 1995:3-10.
17. Cardinal BJ, Spaziani MD. ADA compliance and the accessibility of physical activity facilities in Western Oregon. *Am J Health Promot.* 2003;17:197-201.
18. Fioni SF, McClain L, Degnan JM, et al. Accessibility of physical fitness facilities in the Kansas City metropolitan area. *Top Spinal Cord Inj Rehabil.* 1998;3:66-78.
19. North Carolina Office on Disability and Health. *Removing Barriers to Health Clubs and Fitness Facilities.* Chapel Hill, NC: Frank Porter Graham Child Development Center; 2001.
20. Brownson RC, Baker EA, Housemann RA, et al. Environmental and policy determinants of physical activity in the United States. *Am J Public Health.* 2001;91:1995-2003.
21. Kahn EB, Ramsey LT, Brownson RC, et al. The effectiveness of interventions to increase physical activity. A systematic review. *Am J Prev Med.* 2002;22(4 suppl):73-107.
22. Pikora TJ, Bull FCL, Jamrozik K, et al. Developing a reliable audit instrument to measure the physical environment for physical activity. *Am J Prev Med.* 2003;23:187-194.
23. Nary DE, Froehlich AK, White GW. Accessibility of fitness facilities for persons with disabilities using wheelchairs. *Top Spinal Cord Inj Rehabil.* 2000;6:87-98.
24. West P. Barriers to participation in park and recreation areas by the physically and mentally impaired. *Parks Rec.* 1985;20:67.
25. Lathrop D. Well, well, well. The challenge of keeping fit after rehab is taken on by those who need it. *Rehab Manage.* 1996;9:58-59.
26. Rimmer JH, Riley B, Wang E, Rauworth A. Development and validation of AIMFREE: Accessibility Instruments Measuring Fitness and Recreation Environments. *Disabil Rehabil.* 2004;26:1087-1095.
27. Rimmer JH, Nicola T, Riley B, Creviston T. Exercise training for African Americans with disabilities residing in difficult social environments. *Am J Prev Med.* 2002;23:290-295.
28. Coyle CP, Santiago MC. Aerobic exercise training and depressive symptomatology in adults with physical disabilities. *Arch Phys Med Rehabil.* 1995;76:647-652.
29. Petajan JH, Gappmaier E, White AT, et al. Impact of aerobic training on fitness and quality of life in multiple sclerosis. *Ann Neurol.* 1996;39:432-441.
30. Santiago MC, Coyle CP, Kinney WB. Aerobic exercise effect on individuals with physical disabilities. *Arch Phys Med Rehabil.* 1993;74:1192-1198.

Health Promotion

A fusion of the best of science and the best of practice — together, to produce the greatest impact.



DIMENSIONS OF OPTIMAL HEALTH

Definition of Health Promotion

“Health Promotion is the science and art of helping people change their lifestyle to move toward a state of optimal health. Optimal health is defined as a balance of physical, emotional, social, spiritual and intellectual health. Lifestyle change can be facilitated through a combination of efforts to enhance awareness, change behavior and create environments that support good health practices. Of the three, supportive environments will probably have the greatest impact in producing lasting change.”

(O'Donnell, *American Journal of Health Promotion*, 1989, 3(3):5.)

“The *American Journal of Health Promotion* provides a forum for that rare commodity — *practical and intellectual exchange between researchers and practitioners.*”

Kenneth E. Warner, PhD

Avedis Donabedian Distinguished University Professor of Public Health School of Public Health, University of Michigan

“The contents of the *American Journal of Health Promotion* are *timely, relevant*, and most important, *written and reviewed by the most respected researchers in our field.*”

David R. Anderson, PhD

Vice Programs and Technology, StayWell Health Management

Stay on top of the science and art of health promotion with your own subscription to the *American Journal of Health Promotion*.

Subscribe today...

ANNUAL SUBSCRIPTION RATES: (Good through 12/31/05)

	Individual	Institution
U.S.	\$99.95	\$144.85
Canada and Mexico	\$108.95	\$153.85
Other Countries	\$117.95	\$162.95

CALL 800-783-9913 (U.S. ONLY) or 818-760-8520

OR FIND US ON THE WEB AT

<http://www.HealthPromotionJournal.com>

Editor in Chief

Michael P. O'Donnell, PhD, MBA, MPH

Associate Editors in Chief

Bradley J. Cardinal, PhD

Diane H. Morris, PhD, RD

Judy D. Sheeska, PhD, RD

Mark G. Wilson, HSD

SECTION EDITORS

Interventions

Fitness

Barry A. Franklin, PhD

Medical Self-Care

Donald M. Vickery, MD

Nutrition

Karen Glanz, PhD, MPH

Smoking Control

Michael P. Eriksen, ScD

Weight Control

Kelly D. Brownell, PhD

Stress Management

Cary Cooper, CBE

Mind-Body Health

Kenneth R. Pelletier, PhD, MD (hc)

Social Health

Kenneth R. McLeroy, PhD

Spiritual Health

Larry S. Chapman, MPH

Strategies

Behavior Change

James F. Prochaska, PhD

Culture Change

Daniel Stokols, PhD

Health Policy

Kenneth E. Warner, PhD

Applications

Underserved Populations

Ronald L. Braithwaite, PhD

Health Promoting Community Design

Jo Anne L. Earp, ScD

Research

Data Base

David R. Anderson, PhD

Financial Analysis

Ron Z. Goetzel, PhD

Method, Issues, and Results in Evaluation and Research

Lawrence W. Green, DrPH

Qualitative Research

Marjorie MacDonald, BN, PhD

Measurement Issues

Shawna L. Mercer, MSc, PhD

The Art of Health Promotion

Larry S. Chapman, MPH

